



AU9870091

(12) PATENT ABSTRACT (11) Document No. AU-A-70091/98
(19) AUSTRALIAN PATENT OFFICE

- (54) Title
INDUCING ANTIBODY RESPONSE AGAINST SELF-PROTEINS WITH THE AID OF FOREIGN T-CELL
EPITOPES
- (51)^s International Patent Classification(s)
A61K 039/00 A61K 039/385
- (21) Application No. : 70091/98 (22) Application Date : 11/06/98
- (30) Priority Data
- (31) Number (32) Date (33) Country
964/93 26/08/93 DK DENMARK
- (43) Publication Date : 30/07/98
- (62) Related to Division(s) : 76080/94
- (71) Applicant(s)
MOURITSEN & ELSNER A/S
- (72) Inventor(s)
SOREN MOURITSEN; HENRIK ELSNER
- (74) Attorney or Agent
CARTER SMITH & BEADLE, Qantas House, 2 Railway Parade, CAMBERWELL VIC 3124
- (57)

A novel method for utilizing the immune apparatus to remove and/or down-regulate self-proteins consists in inserting one or more foreign T-cell epitopes in such proteins by molecular biological means, thereby rendering said proteins immunogenic. The modulated self-proteins can be used as autovaccines against undesirable proteins in humans or animals, said autovaccine being useful as vaccines against a number of diseases, e.g. cancer, chronic inflammatory diseases, rheumatoid arthritis, inflammatory bowel diseases, allergic symptoms or diabetes mellitus.

Claim

1. A method for the modification of self-proteins so as to induce antibody response against the unmodified self-proteins following administration of said modified self-proteins to the host, characterized by providing a self-protein analog by molecular biological means by substitution of one or more peptide fragments of the self-protein by a corresponding number of peptides known to contain immunodominant foreign T-cell epitopes, said substitution being carried out so as to essentially preserve the overall tertiary structure of the original self-protein.

2. A method according to claim 1, wherein said immunodominant foreign T-cell epitopes are inserted so as to preserve flanking regions from the original self-protein on both sides of the T-cell epitope.

4. An autovaccine against undesirable self-proteins in humans or animals, characterized in that it comprises one or more self-protein analogs modified according to any of claims 1-3 and formulated with pharmaceutically acceptable adjuvants, such as calcium phosphate, saponin, quil A and biodegradable polymers.

ABSTRACT

A novel method for utilizing the immune apparatus to remove and/or down-regulate self-proteins consists in inserting one or more foreign T-cell epitopes in such proteins by molecular biological means, thereby rendering said proteins immunogenic. The modulated self-proteins can be used as autovaccines against undesirable proteins in humans or animals, said autovaccine being useful as vaccines against a number of diseases, e.g. cancer, chronic inflammatory diseases, rheumatoid arthritis, inflammatory bowel diseases, allergic symptoms or diabetes mellitus.

11
12
13
14
15

16
17

18
19

20
21

AUSTRALIA

Patents Act 1990

COMPLETE SPECIFICATION

FOR A STANDARD PATENT

ORIGINAL

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199

INDUCING ANTIBODY RESPONSE AGAINST SELF-PROTEINS WITH THE
AID OF FOREIGN T-CELL EPITOPES

5 Field of the invention

This invention concerns a novel method for utilizing the immune apparatus to remove and/or down-regulate self-proteins, the presence of which somehow is unwanted in
10 the individual. These could be proteins which are causing disease and/or other undesirable symptoms or signs of disease. Such proteins are removed by circulating autoantibodies which specifically are induced by vaccination. This invention describes a method for developing such
15 autovaccines.

Background of the invention

Physiologically, the vertebrate immune system serves as a
20 defense mechanism against invasion of the body by infectious objects such as micro-organisms. Foreign proteins are effectively removed via the reticuloendothelial system by highly specific circulating antibodies, and viruses and bacteria are attacked by a complex battery of
25 cellular and humoral mechanisms including antibodies, cytotoxic T lymphocytes, Natural Killer cells, complement etc. The leader of this battle is the T helper (T_H) lymphocyte which, in collaboration with the Antigen Presenting Cells (APC), regulate the immune defense via a complex
30 network of cytokines.

T_H lymphocytes recognize protein antigens presented on the surface of the APC. They do not recognize, however, native antigen per se. Instead, they appear to recognize
35 a complex ligand consisting of two components, a "processed" (fragmented) protein antigen (the so-called T

cell epitope) and a Major Histocompatibility Complex class II molecule (O. Werdelin et al., Imm. Rev. 106, 181 (1988)). This recognition eventually enables the T_H lymphocyte specifically to help B lymphocytes to produce specific antibodies towards the intact protein antigen (Werdelin et al., supra). A given T cell only recognizes a certain antigen-MHC combination and will not recognize the same or another antigen presented by a gene product of another MHC allele. This phenomenon is called MHC restriction.

Normally the individual's own proteins (the so-called self- or autoproteins) are not attacked by the immune apparatus. The described events thus generally are beneficial to the individual, but in rare cases the process goes wrong, and the immune system turns towards the individual's own components, which may lead to an autoimmune disease.

The presence of some self-proteins is inexpedient in situations where they, in elevated levels, induce disease symptoms. High levels of immunoglobulins of the IgE class are e.g. known to be important for the induction of type I allergy, and tumor necrosis factor I (TNF α) is known to be able to cause cachexia in cancer patients and patients suffering from other chronic diseases (H.N. Langstein et al., Cancer Res. 51, 2302-2306, 1991). TNF α also plays important roles in the inflammatory process (W.P. Arend et al., Arthritis Rheum. 33, 305-315, 1990). Hormones in sex-hormone dependent cancer are other examples of proteins which are unwanted in certain situations. There is therefore a need for the provision of a method for the development of autovaccines against such self-proteins.

Fragments of self-proteins are also presented by the APC, but normally such fragments are ignored or not recognized by the T helper lymphocytes. This is the main reason why individuals generally do not harbor autoantibodies in their serum.

It is, however, possible artificially to induce antibodies against self-proteins. This can be done, as previously mentioned, by covalent conjugation of the self-protein to an appropriate large foreign carrier protein as e.g. tetanus toxoid or key-hole limpet hemocyanin (KLH). Talwar et al. (G.P. Talwar et al, Int. J. Immunopharmacol. 14, 511-514, 1992) have been able to prevent reproduction in women using a vaccine consisting of a conjugate of human chorionic gonadotropin and tetanus toxoid. There are also other examples of such autoimmunogenic conjugates which have been used therapeutically in man and in animal models (D.R. Stanworth et al., Lancet 336, 1279-1281 (1990)). During the processing of such conjugates in the APC, the necessary T_H lymphocyte stimulatory epitopes are provided from the foreign protein eventually leading to the induction of antibodies against the self-protein as well as against the carrier protein. One disadvantage of using this principle is, however, that the antibody response towards the self-protein will be limited due to shielding of epitopes by the covalently linked carrier protein. Another disadvantage is the increased risk of inducing allergic side-effects due to the contemporary induction of an undesired very strong antibody response against the foreign carrier protein.

Other researchers have conjugated a single peptide predicted to be a T cell epitope chemically as a carrier to a self-peptide [D-Lys⁶]GnRH which is a decapeptide acting as a hapten and managed to induce an autoantibody response with MHC restriction to that particular T cell

epitope (S. Sad et al., Immunology 76, 599-603, 1992). This method seems to be more effective compared with conjugation to large carrier proteins. However, it will only induce antibodies in a population expressing the appropriate MHC molecules. This means that a rather large number of different T cell epitopes has to be conjugated to the self-peptide which will eventually disturb the B cell epitopes on the surface of the self-peptide. Extensive conjugation of proteins may furthermore have the opposite effect with regard to immunogenicity (international patent application No. WO 87/00056) and the surface exposed peptide T cell epitopes may be destroyed by proteolytic enzymes during antigen processing (S. Mouritsen, Scand. J. Immunol. 30, 723, 1989), making that method inexpedient. Also, the exact structure of such multi-conjugated self-peptides will not be chemically and pharmaceutically well-defined.

Recently an improved method has been proposed for breaking the B cell autotolerance by chemical conjugation of B and optionally also peptide T cell epitopes to a high molecular weight dextran molecule (WO 93/23076 published November 25, 1993). The disadvantages mentioned above, however, also hold true for said method.

It has been suggested previously that a universally recognized strong T cell epitope could be associated with a foreign peptide having an antigenic structure representing a B-cell epitope using recombinant DNA technology (EP-A2-0 343 460) It has also been suggested to use peptidyl resin conjugates comprising an immunogenic or antigenic peptide incorporating a helper T-cell (T_H lymphocyte) reactive epitope and preferably a B-cell reactive epitope in the preparation of immunogenic compositions, e.g. vaccines. The conjugates are prepared by solid phase synthesis, preferably on a polyamide resin. (WO

90/15627). While the intent is to increase an antibody response towards the peptides in question, it has not been proposed that this could be done with the purpose of breaking the autotolerance of the immune system and induce an antibody response against self-proteins. Using these methods for induction of autoantibodies against self-proteins one a priori would expect the same rules to be true with regard to the above-mentioned limitations of the MHC restriction of the response. Surprisingly, however, by modulation of self-proteins using the method according to the invention, wherein a self-protein analog is produced by substitution of one or more peptide fragments by a corresponding number of peptides known to contain immunodominant T-cell epitopes, said substitution being carried out so as to essentially preserve the overall tertiary structure of the original self-protein, it proved possible to induce an equally fast and even a stronger autoantibody response against TNF α despite the fact that the inserted T cell epitope used was not restricted to the MHC molecules of the immunized mice, vide Example 3 below. The reason for this observation is not clear but may be due to the appearance of new MHC binding segments in the mutagenized area in the self-protein. However, the experiment shown in example 5 demonstrates that this is probably not the case, since synthetic peptides representing overlapping regions of the implanted ovalbumin T cell epitope in ubiquitin did not bind strongly to any of the MHC class II molecules of the H-2^k mice in which this recombinant molecule was highly immunogenic.

Most of the potential MHC class II binding segments of a protein are normally cryptic and will not be presented to the host T cells by the antigen presenting cells (S. Mouritsen et al, Scand. J. Immunol. 34, 421, 1991). The observed lacking correspondence between the MHC restric-

tion of the inserted T cell epitope and the restriction of the antibody response could perhaps be due to a general disturbance of the intra-molecular competition of binding to MHC molecules by different self-protein segments. According to the method of the invention non-tolerized cryptic self-protein segments may be presented to the T cells leading to breaking of the T cell as well as the B cell autotolerance towards the protein. In accordance with the invention and illustrated in all the examples described below, a fragment of the self-protein was substituted with a foreign T cell epitope. This deletion followed by a substitution with an other protein fragment minimally obscure the tertiary structure of the self-proteins, but may also contribute strongly to the disturbance of said intramolecular competition of the MHC class II binding self-segments. This concept is therefore clearly different from the above-mentioned prior art mechanisms and methods. Independently of the operating mechanism of action by the method according to the invention, it is more technically advantageous compared to the known methods for breaking the B cell autotolerance, since it is possible to induce antibodies in a broad population of MHC molecules by insertion of a minimal number of different foreign T cell epitopes.

The present invention thus is based on the surprising fact that injection of recombinant self-proteins, which have been appropriately modulated by deletion of one or more peptide fragments and simultaneous insertion of a corresponding number of foreign T cell epitopes, so as to produce a self-protein analog with an essentially preserved tertiary structure, induces a profound autoantibody response against the unmodified self-proteins. Surprisingly the MHC-restriction of the auto-antibody response induced was not necessarily confined to that of the inserted T cell epitope. By inducing minimal tertiary

structural changes in the highly conserved self-protein ubiquitin, as well as in TNF α , foreign T cell epitopes having a length of 12-15 amino acids were inserted using genetic engineering methods. These recombinant self-protein analogs were purified, emulsified in adjuvant and injected into mice. Within only one week an autoantibody response against ubiquitin could be detected in serum from these mice. Non-modified, recombinant ubiquitin treated and injected in the same way was not able to induce a response.

By using this principle for developing vaccines against undesirable proteins, the risk of inducing allergic side-effect is reduced, and toxic self-proteins such as TNF α can simultaneously be detoxified by removing or mutating biologically active protein segments. The epitope-shielding effect described above is not a problem, and autoantibodies against ubiquitin were induced much faster as compared to the known technique, in which the self-protein is conjugated to a carrier protein or peptide. Importantly, by this method it furthermore seems possible to temporarily break the autotolerance of the T cells as well as that of the B cells of the individual, and such recombinant proteins will be self-immunogenic in a large population expressing many different MHC class II molecules.

The vaccine according to the invention consists of one or more self-protein analogs modulated as described above and formulated with suitable adjuvants, such as calcium phosphate, saponin, quil A or biodegradable polymers. The modulated self-protein analogs may be prepared as fusion proteins with suitable, immunologically active cytokines, such as GM-CSF or interleukin 2.

Accordingly, in one aspect the invention provides a method for preparing a modified self-protein, said self-protein being derived from a vertebrate, the method comprising substituting, by molecular biological means, one or more peptides in the self-protein with a corresponding number of peptides, wherein

- 5 a) the substituting peptides contain immunodominant T-cell epitopes which are foreign to the vertebrate, and
- b) the substitution(s) carried out essentially preserve(s) the overall tertiary structure of the unmodified self-protein,

and thereafter confirming that said modified self-protein induces, in the vertebrate, an
 10 antibody response against the unmodified self-protein following delivery to the vertebrate of said modified self-protein.

In another aspect, the invention provides an autovaccine against undesirable self-proteins in humans or animals,

15 characterised in that it comprises one or more self-proteins modified according to the foregoing method and formulated with pharmaceutically acceptable adjuvants, such as calcium phosphate, saponin, quil A or biodegradable polymers.

20 In another aspect the invention provides a modified self-protein comprising one or more peptide fragments containing immunodominant foreign T-cell epitopes wherein the modified self-protein, when introduced in a vertebrate wherein the unmodified self-protein naturally occurs, induces production of antibodies which cross-react with the self-protein,

25 with the proviso that the foreign T-cell epitopes(s) is/are inserted so as to preserve flanking regions from the unmodified self-protein on both sides of said T-cell epitope and that the overall tertiary structure of the modified self-protein as compared with the unmodified self-protein essentially being preserved.



MAW:JMD:#23631

18 February 1999

The autovaccine may i.a. be a vaccine against TNF α or γ -interferon for the treatment of patients with cachexia, e.g. cancer patients, or a vaccine against IgE for the treatment of patients with allergy. Furthermore, it may
5 be a vaccine against TNF α , TNF β or interleukin 1 for the treatment of patients with chronic inflammatory diseases.

The invention is illustrated in the following examples:

10 EXAMPLE 1

Substitution of foreign T cell epitopes into ubiquitin

An overview of this procedure is shown in fig. 1 using
15 the T cell epitope MP7 as example. The gene sequences representing MP7 (MP7.1-C and MP7.1-NC) were synthesized as two complementary oligonucleotides designed with appropriate restriction enzyme cloning sites. The amino acid sequence of MP7 is PELFEALQKLFKHAY, Mouritsen et al,
20 Scand. J. Immunol. 30, 723-730, 1989. The oligonucleotides were synthesized using conventional, automatic solid phase oligonucleotide synthesis and purified using agarose gel electrophoresis and low melting agarose. The desired bands were cut out from the gels, and known quantities of oligonucleotides were mixed, heated to 5C below
25 their theoretical melting point (usually to approximately 65C) for 1-2 hours, and slowly cooled to 37C. At this temperature the hybridized oligonucleotides were ligated to the vector fragments containing part of the ubiquitin
30 gene. The subsequent analysis of positive clones using restriction fragment analysis and DNA sequencing was done by conventional methods ("Molecular Cloning", Eds.: T. Maniatis et al. 2 ed. CSH Laboratory Press, 1989).

EXAMPLE 2Induction of autoantibodies against ubiquitin by vaccination with modified ubiquitin analogs

5 Genes containing sequences encoding the foreign T cell epitopes OVA (325-336) from ovalbumin (OVA) and HEL (50-61) from hen eggwhite lysozyme (HEL), respectively, were expressed in E. coli strain, AR58 under control of the heat sensitive S repressor regulated promotor. Expression of the recombinant ubiquitin proteins were verified using a polyclonal anti-ubiquitin antibody and Western-blotting ("Antibodies", Eds.: D. Harlow et al., CSH Laboratory Press, 1988). The recombinant proteins were purified using conventional methods (Maniatis et al., supra).

Mice were injected i.p. with 100 µg of ubiquitin or its analogs in phosphate buffered saline (PBS) emulsified in Freund's Complete adjuvant. Booster injections of the same amount of antigen emulsified 1:1 in Freund's Incomplete adjuvant were performed i.p. at days 14 and 28. Five Balb/c mice in each group were examined and blood samples were examined for the presence of anti-ubiquitin antibodies on day 7, 14, 21, 28, 35, and 42 using conventional ELISA methodology.

The results exemplified by the antibody response against two different ubiquitin analogs containing the T cell epitopes OVA(325-336) and HEL(50-61), respectively, are shown in fig. 2. The amino acid sequence of the inserted OVA(325-336) epitope is: QAVHAAHAEINE and the amino acid sequence of the HEL(50-61) epitope is STDYGILQINSR.

A clear antibody response against native ubiquitin could be detected within only one week from the first injection of antigen reaching a maximum within 2 weeks. Anti-

ubiquitin antibodies produced in rabbits by covalently conjugating ubiquitin to bovine immunoglobulin reached maximum values after a much longer immunization period (data not shown).

5

EXAMPLE 3

Induction of autoantibodies against tumor necrosis factors (TNF) by vaccination with appropriately modified TNF analogs

10

The gene coding for the structural part of the native murine TNF α protein (MR101) was obtained by Polymerase Chain Reaction (PCR) cloning of the DNA. In the MR103 TNF α analog the ovalbumin (OVA) H-2^d restricted T cell epitope sequence 325-334 (QAVHAAHAET) replaces the amino acids 26-35 in the cloned TNF α sequence, a substitution of an amphipathic α -helix. Substitutions in this region of the TNF α detoxifies the recombinant protein (X. Van Ostade et al., Nature 361, 266-269, 1993). In the MR105 analog the H-2^k restricted T cell epitope from hen eggwhite lysozyme (HEL), amino acid sequence 81-96 (SALLSSDITASVNCAK) replaces the amino acids 5-20 in the cloned TNF α sequence. In the MR106 TNF α mutant the same epitope, amino acid sequence 81-95 (SALLSSDITASVNCA) replaces the amino acids 126-140 in the cloned TNF α sequence. The genetic constructions are shown in Fig. 3. Different techniques compared to the technique described in example 1 were used for exchanging parts of the TNF α gene with DNA coding for T cell epitopes. The MR105 and 106 constructs were made by introducing the mutant sequence by PCR recloning a part of the TNF α gene flanking the intended site for introducing the T cell epitope. The mutant oligonucleotide primer contained both a DNA sequence homologous to the TNF α DNA sequence as well as a DNA sequence encoding the T cell epitope. The PCR re-

35

cloned part of the TNF α gene was subsequently cut with appropriate restriction enzymes and cloned into the MR101 gene. The MR103 construction was made by a modification of the "splicing by overlap extension" PCR technique (R. M. Horton et al., Gene 77, 61, 1989). Here two PCR products are produced, each covering a part of the TNF α gene, and additionally each PCR product contains half of the T cell epitope sequence. The complete mutant TNF α gene was subsequently made by combining the two PCR products in a second PCR. Finally, the complete genetic constructions were inserted into protein expression vectors. Subsequently, all genetic constructions were analyzed by restriction fragment analysis and DNA sequencing using conventional methods ("Molecular Cloning", Eds.: T. Maniatis et al. 2.ed. CSH Laboratory Press, 1989). The recombinant proteins were expressed in E.coli and purified by conventional protein purification methods.

Groups of BALB/c (MHC haplotype H-2^d) and C3H (MHC haplotype H-2^k) mice, respectively, were immunized subcutaneously with 100 Tg of semi-purified MR103 and MR106 emulsified in Freund's complete adjuvant. Every second week the immunizations were repeated using incomplete Freund's adjuvant. All mice developed an early and strong antibody response against biologically active MR101. This was measured by a direct ELISA method using passively adsorbed 100% pure MR101 (Fig. 4). Control mice immunized with MR101 and PBS, respectively, showed no antibody reactivity towards MR101.

Strikingly, the antibody response towards MR101 was not MHC restricted corresponding to the implanted T cell epitopes, since both mice strains of different MHC haplotypes responded well to MR103 and MR106 containing differently restricted T-cell epitopes (Fig. 4). Taken together these results illustrate (a) the ability of the

self-protein analogs produced by the method according to the invention to induce autoantibodies towards a secreted autoprotein and (b) the improved efficiency of the herein described method with regard to inducing a response in a broader MHC population than predicted by the MHC binding ability of the inserted T cell epitopes. The immune response against MR101 induced by the recombinant self-protein analogs MR103 and MR106 was much stronger and more high-titered compared to the immune response induced by aldehyde conjugated MR101 (see Example 4).

EXAMPLE 4

Induction of autoantibodies against TNF α by self-protein analogs produced by the method according to the invention compared to unmodified self-protein conjugated to E. coli proteins.

The induction of autoantibodies against TNF α by the method of the present invention has been directly compared to the autoantibody response induced when using a conjugate of TNF α and E. coli proteins, which must contain small single T cell epitope peptides as well as larger foreign carrier proteins.

Semi-purified recombinant murine TNF α (MR101) was conjugated to E. coli proteins in PBS, pH 7.4, using 0.5% formaldehyde. Conjugation of the proteins was confirmed by SDS-PAGE. These conjugates were subsequently used for immunization of C3H mice. Another group of C3H mice was vaccinated with semi-purified non-conjugated self-protein analog MR105. About 100 μ g of recombinant TNF α analog and conjugate were emulsified 1:1 in Freund's complete adjuvant and injected subcutaneously in each group of mice. MR105 is biologically inactive as judged by the L929 bioassay for TNF α . In subsequent immunizations every second

week incomplete Freund's adjuvant was used. Both groups eventually developed autoantibodies against highly purified biologically active MR101 as determined by ELISA, but the immune response against the non-conjugated analog MR105 produced by the method of the invention was induced earlier and was of a higher titer (Fig. 5).

EXAMPLE 5

10 The possible MHC class II binding of peptides representing overlapping sequences of self-protein as well as of the ovalbumin T cell epitope inserted in ubiquitin.

Peptide-MHC complexes were obtained by incubating ^{125}I -labeled peptide (10-100 nM) with affinity purified MHC class II molecules (2-10 μM) at room temperature for 3 days (S. Mouritsen, J. Immunol. 148, 1438-1444, 1992). The following peptides were used as radiolabeled markers of binding: Hb(54-76)Y which binds strongly to the E^{K} molecule and HEL(46-61)Y which binds strongly to the A^{K} molecule. These complexes were co-incubated with large amounts of cold (non-radiolabeled) peptide (> 550 Tm) which is sufficient to inhibit totally all immunologically relevant MHC class II binding. Either the same peptides were used, or three different overlapping peptides were used, said peptides representing the flanking regions as well as the entire OVA(325-336) T cell epitope which was substituted into ubiquitin (see Example 2). The three peptides were: TITLEVEPSQAVHAA (U(12-26)), PSQAVHAAHAEINEKE (U(19-34)) and HAEINEKEGIPPDQQ (U(27-41)). The reaction buffer contained 8 mM citrate, 17 mM phosphate, and 0.05% NP-40 (pH 5) and peptide-MHC class II complexes were separated (in duplicate) from free peptide by gel filtration using G25 spun columns. Both the radioactivities of the excluded "void" volume and of the included volume were measured by gamma spectrometry. The

competitive inhibition of maximal binding (in percent) by addition of cold peptide was calculated. The results are shown in Table I.

5 TABLE I.

Peptide/		Hb(64-76)	HEL(46-61)	U(12-26)	U(19-34)	U(27-41)
MHC						
10	A ^k	28.6	<u>97.4</u>	35.3	44.6	7.8
	E ^k	<u>92.6</u>	0.0	45.6	12.2	0.0

It can be seen that total inhibition of the binding of the radiolabeled peptides Hb(64-76)Y and HEL(46-61)Y to E^k and A^k respectively could only be achieved using cold versions of the same peptides. Although some inhibition of binding was seen by U(12-26) and U(19-34) using these extreme amounts of cold peptide, it is likely that the affinity of these peptides to the H-2^k MHC class II molecules is very low. Therefore this seems not to be sufficient to explain the strong immunogenicity in the H-2^k mouse strain of the ubiquitin analog containing the ovalbumin T cell epitope. More likely, other and previously non-tolerized self-epitopes are presented to the T cell in these animals.

EXAMPLE 6

30 Treatment of diabetes or inflammatory disease by vaccination with appropriately modified TNF α analogs

Genes coding for TNF α are modified by insertion of appropriate gene segments coding for T cell epitopes derived from e.g. tetanus toxin or influenza hemagglutinin. Such genes are expressed in appropriate expression vectors in

e.g. *E. coli* or insect cells. The recombinant TNF α proteins were purified using conventional methods ("Molecular Cloning", Eds.: T. Maniatis et al. 2. ed. CSH Laboratory Press, 1989).

5

Optionally such recombinant proteins can be coupled to immunologically active cytokines such as GM-CSF or interleukin 2.

10

The recombinant proteins can be formulated with appropriate adjuvants and administered as an anti-TNF α vaccine to patients suffering from diseases where TNF α is important for the pathogenesis. The induced anti-TNF α antibodies will thereby affect the diseases.

15

One example of said diseases is the chronic inflammatory diseases such as e.g. rheumatoid arthritis where TNF α is believed to play an important role (reviewed in: F.M. Brennan et al., Br. J. Rheumatol. 31, 293-298, 1992).

20

TNF α is also believed to play an important role in the cachectic conditions seen in cancer and in chronic infectious diseases such as AIDS (reviewed in M. Odeh. J. Intern. Med. 228, 549-556, 1990). It is also known that TNF participates in septic shock (reviewed in: B.P. Giroir, Crit. Care. Med., 21, 780-789, 1993). Furthermore, it has been shown that TNF α may play a pathogenetic role in the development of type II diabetes mellitus (CH Lang et al., Endocrinology 130, 43-52, 1992).

25

30 LEGENDS TO FIGURES

Fig. 1. Schematic overview of the cloning strategy used in the construction of a ubiquitin gene with an implanted foreign T cell epitope (MP7). Restriction enzyme digestions, hybridization and ligation procedures are indi-

35

cated with arrows. Fragment sizes are shown in parentheses.

Fig. 2. Reactivity toward immobilized bovine ubiquitin
5 in sera from mice immunized with recombinant ubiquitin
and analogs containing the implanted T cell epitopes
OVA(325-336) and HEL (50-61), respectively. Fig. 2a) sera
from Balb/c mice immunized with recombinant ubiquitin
containing OVA(325-336). Fig. 2b) sera from Balb/c mice
10 immunized with recombinant ubiquitin containing the T
cell epitope HEL(50-61). Fig. 2c) sera from Balb/c mice
immunized with recombinant non-modified ubiquitin. Sera
(diluted 1:100) were tested in a standard ELISA assay us-
ing non-modified bovine ubiquitin immobilized on the
15 solid phase.

Fig. 3. Schematic overview of the cloning strategy used
in the construction of the recombinant TNF α mutants. PCR
products and restriction enzyme digestions are indicated.
20

Fig. 4. Induction of TNF α autoantibodies by vaccination
of Balb/c or C3H mice with semipurified MR103 and MR106.
The antibody titers were measured by ELISA and expressed
as arbitrary units (AU) referring to a strong standard
25 anti-serum from one mouse. The plotted values represent a
mean titer for 5 animals. Freund's complete adjuvant was
used as adjuvant for the first immunization. All subse-
quent immunizations at 14 days intervals were done with
Freund's incomplete adjuvant. Mice immunized in parallel
30 with native MR101 in PBS did not develop detectable TNF α
autoantibodies (data not shown). Non-detectable antibody
titers were assigned the titer value 1.

Fig. 5. Anti TNF α autoantibodies induced by vaccination
35 with non-conjugated MR105 and MR101 conjugated to E. coli
proteins, respectively. C3H mice and Balb/c mice were im-

munized with both preparations. The immunizations, measurements and calculations of mean antibody titers were done as described in example 4.

1957
1958
1959
1960

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for preparing a modified self-protein, said self-protein being derived from a vertebrate, the method comprising substituting, by molecular biological means, one or more peptides in the self-protein with a corresponding number of peptides, wherein
 - a) the substituting peptides contain immunodominant T-cell epitopes which are foreign to the vertebrate, and
 - b) the substitution(s) carried out essentially preserve(s) the overall tertiary structure of the unmodified self-protein,
 and thereafter confirming that said modified self-protein induces, in the vertebrate, an antibody response against the unmodified self-protein following delivery to the vertebrate of said modified self-protein.
2. A method according to claim 1, wherein said immunodominant foreign T-cell epitopes are inserted so as to preserve flanking regions from the unmodified self-protein on both sides of the T-cell epitope.
3. A method according to any of claims 1 or 2, wherein the immunodominant T-cell epitope(s) originate(s) from tetanus toxoid or diphtheria toxoid.
4. An autovaccine against undesirable self-proteins in humans or animals, characterised in that it comprises one or more self-proteins modified according to any of claims 1-3 and formulated with pharmaceutically acceptable adjuvants, such as calcium phosphate, saponin, quil A and biodegradable polymers.
5. An autovaccine according to claim 4, characterised in that the modified self-protein is present in the form of a fusion protein with suitable, immunologically active cytokines, such as GM-CSF or interleukin 2.
6. An autovaccine according to claim 4, characterised in that it is a vaccine against TNF α or γ -interferon for the treatment of patients susceptible to cachexia, e.g. cancer patients.
7. An autovaccine according to claim 4, characterised in that it is a vaccine against IgE for the treatment of patients with allergy.



8. An autovaccine according to claim 4, characterised in that it is a vaccine against $\text{TNF}\alpha$, $\text{TNF}\beta$ or interleukin 1 for the treatment of patients with chronic inflammatory diseases.

5 9. An autovaccine according to claim 8, characterised in that it is a vaccine for treatment of patients with rheumatoid arthritis or an inflammatory bowel disease.

10 10. An autovaccine according to claim 4 or 5, characterised in that it is a vaccine against $\text{TNF}\alpha$ for the treatment of diabetes mellitus.

11. A method according to any one of claims 1 to 3 substantially as hereinbefore described.

15 12. An autovaccine according to any one of claims 4 to 10 substantially as hereinbefore described.

13. A method of inducing an antibody response against a self-protein, said self-protein being derived from a vertebrate, the method comprising:

20 introducing, in the vertebrate, an effective amount of the self-protein which has been modified by substitution of one or more peptides in the self-protein with a corresponding number of peptides comprising immunodominant T-cell epitopes which are foreign to the vertebrate while preserving the overall tertiary structure of the self-protein.

25 14. Use of a modified self-protein modified according to the method of any one of claims 1-3 for inducing antibodies against the unmodified self-protein in a vertebrate harbouring the self-protein.

30 15. A modified self-protein comprising one or more peptide fragments containing immunodominant foreign T-cell epitopes wherein the modified self-protein, when introduced in a vertebrate wherein the unmodified self-protein naturally occurs, induces production of antibodies which cross-react with the self-protein,

35 with the proviso that the foreign T-cell epitopes(s) is/are inserted so as to preserve flanking regions from the unmodified self-protein on both sides of said T-



cell epitope and that the overall tertiary structure of the modified self-protein as compared with the unmodified self-protein essentially being preserved.

16. A modified self-protein according to claim 15 formulated with
5 pharmaceutically acceptable adjuvants, such as calcium phosphate, saponin, quil A and biodegradable polymers.

17. A modified self-protein according to claim 15 in the form of a fusion
10 protein with suitable, immunologically active cytokines, such as GM-CSF or interleukin 2.

18. A modified self-protein according to claim 15 being a protein for use in the treatment of patients susceptible to cachexia, e.g. cancer patients.

19. A modified self-protein according to claim 15 for use in the treatment
15 of patients with allergy.

20. A modified self-protein according to claim 15 for use in the treatment
20 of patients with chronic inflammatory diseases.

21. A modified self-protein according to claim 15 for use in the treatment
of patients with rheumatoid arthritis or an inflammatory bowel disease.

22. A modified self-protein according to claim 15 for use in the treatment
25 of diabetes mellitus.

23. A method according to claim 13 substantially as hereinbefore described.

24. A modified self protein according to any one of claims 15 to 22
30 substantially as hereinbefore described.

DATED: 18 February 1999

CARTER SMITH & BEADLE
Patent Attorneys for the Applicant:

MOURITSEN & ELSNER S/A






FIG. 1

11 05 98 70001

2/5




FIG. 2

Cloning strategy for murine TNF α mutants.




FIG. 3


Anti TNF α auto-antibodies

FIG. 4

11 05 99 10031

